arXiv:math/9811131v2 [math.AG] 15 Jan 2000

Syzygies of Veronese embeddings

Giorgio Ottaviani and Raffaella Paoletti*

Abstract - *We prove that the Veronese embedding* $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}\colon \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ *with* $n \geq 2$, $d \geq 3$ *does not satisfy property* N_p *(according to Green and Lazarsfeld) if* $p \geq 3d-2$. *We make the conjecture that also the converse holds. This is true for* $n = 2$ *and for* $n = d = 3$.

Introduction

Let $\mathbb{P}^n$ be the projective n-space over an algebraically closed field of characteristic zero and let $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)} : \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ be the Veronese embedding associated to the complete linear system $|\ \mathcal{O}_{\mathbb{P}^n}(d)|$. In order to understand the homogeneous ideal $\mathcal{I}$ of $\mathbb{P}^n$ in $\mathbb{P}^N$ as well as its syzygies, it is useful to study some properties about a minimal free resolution of $\mathcal{I}$.
M. Green and R. Lazarsfeld (see [G2], [GL]) introduced the property N_p (Definition 1.3) for a complete projective non singular variety $X \hookrightarrow \mathbb{P}^N$ embedded in $\mathbb{P}^N$ with an ample line bundle L. When property N_p holds for every integer p the resolution of $\mathcal{I}$ is "as nice as possible". M. Green proves in [G2], Theorem 2.2, that $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ satisfies N_p if $p \leq d$. L. Manivel has generalized this result to flag manifolds [M]. The rational normal curves (which are the Veronese embeddings of $\mathbb{P}^1$) satisfy N_p $\forall p$. C. Ciliberto showed us that the results of [G1] imply that $\varphi_{\mathcal{O}_{\mathbb{P}^2}(d)}$ with $d \geq 3$ satisfies N_p if $p \leq 3d-3$. This sufficient condition has been found also by C. Birkenhake in [B1] as a corollary of a more general result. Here we prove that this condition is also necessary (Theorem 3.1) and we formulate (for $n \geq 2$) the following

* both authors were partially supported by GNSAGA and by Progetto Strategico "Calcolo Simbolico" of CNR.

Conjecture $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ *satisfies* $N_p \iff \begin{cases} n=2, d=2, \ \forall p \\ n\geq 3, d=2, \ p\leq 5 \\ n\geq 2, d\geq 3, \ p\leq 3d-3 \end{cases}$.

Our precise result is the following:

Theorem *The implication '$\Longrightarrow$' of the previous conjecture is true.*

Moreover we remark that the implication '$\Longleftarrow$' of the previous conjecture is true in the cases $n=2$ ([G1]), $n=d=3$ ([G1]), $d=2$ ([JPW]). This solves the Problem 4.5 of [EL] (raised by Fulton) in the first cases given by the projective plane and by the cubic embedding of the projective 3-dimensional space.

We remark also that our conjecture would be overcomed by the knowledge of the minimal resolution of the Veronese variety. This is stated as an open problem in [G2] (remark of section 2). Our results can be seen as a step towards this problem.

The paper is organized as follows: in *section* 1 we recall some definitions we will need later and we improve a known cohomological criterion for the property N_p; in *section* 2 we prove our main results and in *section* 3 we fit our results into the literature.

We thank the referee for the improvements suggested in the presentation of the paper.

KEYWORDS: Veronese embedding, free resolution, syzygy.
SUBJECT CLASSIFICATION: 14M20, 14F17.

1 - Notations and preliminaries

Let V be a vector space of dimension $n+1$ over an algebraically closed field $I\!K$ of characteristic 0 and let $\mathbb{P}^n = \mathbb{P}(V^*)$ the projective space associated to the dual space of V.
Note that $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d)) \cong S^dV \quad \forall d\geq 0$.
For any vector bundle E over $\mathbb{P}^n$, we will denote by $H^i(E)$ the i-th cohomology group of E over $\mathbb{P}^n$ and by $E(t)$ the tensor product $E\otimes\mathcal{O}_{\mathbb{P}^n}(t)$.

The following bundles will play a fundamental role in this paper:

Definition 1.1 For any positive integer d, the line bundle $\mathcal{O}_{\mathbb{P}^n}(d)$ is generated by global sections $H^0(\mathcal{O}_{\mathbb{P}^n}(d)) \cong S^dV$ so that the evaluation map $ev : S^dV \otimes \mathcal{O}_{\mathbb{P}^n} \longrightarrow \mathcal{O}_{\mathbb{P}^n}(d)$ is surjective. Call E_d the kernel. Thus the vector bundle E_d is defined by the exact sequence:

$$0 \longrightarrow E_d \longrightarrow S^d V \otimes \mathcal{O}_{\mathbb{P}^n} \xrightarrow{ev} \mathcal{O}_{\mathbb{P}^n}(d) \longrightarrow 0\,. \tag{1.2}$$

It follows immediately from the definition that the bundle E_d has rank $N := rk_{E_d} = \binom{n+d}{n} - 1$ and first Chern class $c_1(E_d) = -d$.

Note that, if $d = 1$, (1.2) is the dualized Euler sequence so that:

$$E_1 \cong \Omega^1_{\mathbb{P}^n}(1) \quad \text{and} \quad \bigwedge^q E_1 \cong \Omega^q_{\mathbb{P}^n}(q)\,.$$

For any integer $d \geq 0$, we will denote by $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ the *Veronese embedding* $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)} : \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ associated to the complete linear system $|\, \mathcal{O}_{\mathbb{P}^n}(d)\,|$ of dimension $N+1 := \binom{n+d}{n}$. Recall that if $[x_0 : \ldots : x_n]$ is a system of homogeneous coordinates on $\mathbb{P}^n$ and $[y_0 : \ldots : y_N]$ on $\mathbb{P}^N = \mathbb{P}\left(H^0\left(\mathcal{O}_{\mathbb{P}^n}(d)\right)^*\right)$, then $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ is the embedding:

$$[x_0 : \ldots : x_n] \hookrightarrow [x_0^d : x_0^{d-1}x_1 : \ldots : x_n^d]\,.$$

With the above notation, let $S := \bigoplus_{k\geq 0} S^k\left(H^0\left(\mathbb{P}^n,\, \mathcal{O}_{\mathbb{P}^n}(d)\right)\right)$ be the homogeneous coordinate ring of $\mathbb{P}^N$ and define the graded S-module $R := \bigoplus_{k\geq 0} H^0\left(\mathbb{P}^n,\, \mathcal{O}_{\mathbb{P}^n}(kd)\right)$. Let

$$0 \longrightarrow \oplus_j S(-j)^{\widetilde{b}_{rj}} \longrightarrow \cdots \longrightarrow \oplus_j S(-j)^{\widetilde{b}_{0j}} \longrightarrow R \longrightarrow 0$$

be a minimal free resolution of R with *graded Betti numbers* $\widetilde{b}_{ij}$.

Definition 1.3 For any integer $p \geq 0$ the embedding $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)} : \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ is said to satisfy *property* N_p if

$$\widetilde{b}_{0j} = \begin{cases} 1 & \text{if } j = 0 \\ 0 & \text{otherwise} \end{cases} \qquad \text{and} \qquad \widetilde{b}_{ij} = 0 \quad \text{for } j \neq i+1 \text{ when } 1 \leq i \leq p\,.$$

Thus:

N_0 means that $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}(\mathbb{P}^n)$ is projectively normal in $\mathbb{P}^N$;

N_1 means that N_0 holds and the ideal I of $\mathbb{P}^n \hookrightarrow \mathbb{P}^N$ is generated by quadrics;

N_2 means that, moreover, the module of syzygies among quadratic generators $Q_i \in I$ is spanned by the relations of the form $\sum L_i Q_i = 0$ where the L_i are *linear* polynomials;

and so on.

Remark 1.4 Let $\mathcal{C} \hookrightarrow \mathbb{P}^d$ be the rational normal curve (of degree d) in $\mathbb{P}^d$. If V is a vector space of dimension 2, then $\mathcal{C} \cong \mathbb{P}(V^*) \hookrightarrow \mathbb{P}^d = \mathbb{P}(S^d V^*)$ is the image of the Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^1}(d)} : \mathbb{P}^1 \hookrightarrow \mathbb{P}^d$.

It is well known (e.g. by using the Eagon-Northcott complex) that the sheaf ideal $\mathcal{I}$ of $\mathcal{C}$ in $\mathcal{O}_{\mathbb{P}^d}$ has the following resolution:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^d}(-d)^{\oplus b_d} \longrightarrow \mathcal{O}_{\mathbb{P}^d}(-d+1)^{\oplus b_{d-1}} \longrightarrow \cdots \longrightarrow \mathcal{O}_{\mathbb{P}^d}(-2)^{\oplus b_2} \longrightarrow \mathcal{I} \longrightarrow 0$$

where $b_k := (k-1)\binom{d}{k}$. So the Veronese embeddings of $\mathbb{P}^1$ satisfy $N_p \ \forall p$.

From [B2], Remark 2.7, and [G1] we have the following cohomological criterion:

Proposition 1.5 *The Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ satisfies property N_p if and only if*

$$H^1\left(\bigwedge^q E_d(jd)\right) = 0 \qquad \text{for} \quad 1 \le q \le p+1 \quad \text{and} \quad \forall j \ge 1\,.$$

$\diamond$

We have the following cohomological criterion, which improves slightly the previous one (in fact $H^2(\bigwedge^q E_d) \simeq H^1\left(\bigwedge^{q-1} E_d(d)\right)$).

Theorem 1.6 *The Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ satisfies property N_p if and only if $H^2(\bigwedge^q E_d) = 0$ for $1 \le q \le p+2$.*

The proof of Theorem 1.6 relies on the following

Proposition 1.7 *If* $H^2(\bigwedge^q E_d) = 0$ *for* $1 \leq q \leq k$, *then* $H^2(\bigwedge^q E_d(t)) = 0$ *for* $1 \leq q \leq k$ *and* $\forall t \geq 0$.

Proof
Consider the two exact sequences:

$$(*) \qquad 0 \longrightarrow \bigwedge^q E_d(t-1) \longrightarrow \bigwedge^q E_d(t) \longrightarrow \bigwedge^q E_d(t)|_{\mathbb{P}^{n-1}} \longrightarrow 0\,,$$

$$(**)\ 0 \longrightarrow \bigwedge^q E_d(t-1) \longrightarrow \bigwedge^q (S^d V) \otimes \mathcal{O}_{\mathbb{P}^n}(t-1) \longrightarrow \bigwedge^{q-1} E_d(t+d-1) \longrightarrow 0\,.$$

The proof is by double induction on n and k. The statement is true for $n = 2$ (Serre duality) and for $k = 1$ (it follows immediately from (1.2)). From the cohomology sequence associated to $(**)$ with $t = 0$ and the inductive hypothesis on k we get $H^3(\bigwedge^q E_d(-1)) = 0$ for $1 \leq q \leq k$. Since $E_d|_{\mathbb{P}^{n-1}} \cong \widetilde{E}_d \oplus \mathcal{O}_{\mathbb{P}^{n-1}}^{\oplus \binom{n+d-1}{n}}$, where $\widetilde{E}_d$ is the vector bundle E_d over $\mathbb{P}^{n-1}$, the previous vanishing implies in the cohomology sequence associated to $(*)$ with $t = 0$ that the hypothesis of the proposition are true on $\mathbb{P}^{n-1}$. Hence by induction on n, $H^2(\mathbb{P}^{n-1}, \bigwedge^q E_d(t)_{|\mathbb{P}^{n-1}}) = 0$ for $1 \leq q \leq k$ and $\forall t \geq 0$. From the cohomology sequence associated to $(*)$ with $q = k$ we get that the map $H^2(\mathbb{P}^n, \bigwedge^k E_d(t-1)) \longrightarrow H^2(\mathbb{P}^n, \bigwedge^k E_d(t))$ is surjective $\forall t \geq 0$ and the thesis follows easily. $\diamond$

Proof of Theorem 1.6
The implication '$\Longrightarrow$' is a consequence of Proposition 1.5. To prove the converse, we may apply Proposition 1.7 and then Proposition 1.5 again.

Proposition 1.8 *If* $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ *satisfies* N_p *then* $\varphi_{\mathcal{O}_{\mathbb{P}^m}(d)}$ *satisfies* N_p $\forall m \leq n$.

Proof
It follows by the remark of section 2 of [G2] (which is an insight into representation theory). $\diamond$

2 - Necessary conditions on property N_p for the Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$

In this section we will prove the following:

Theorem 2.1 *The Veronese embedding* $\varphi_{\mathcal{O}_{\mathbb{P}^n}(d)}$ *does not satisfy* N_{3d-2} *for* $n \geq 2\,,\ d \geq 3$.

Proof
By Proposition 1.8, we can let n=2. By Theorem 1.6 and Serre duality it is enough to show that $H^0\left(\mathbb{P}^2,\, \bigwedge^K E_d(d-3)\right) \neq 0$ with $K := \frac{d(d-3)}{2}$. So the theorem will follow from the following:

Lemma 2.2 *The bundle* $\bigwedge^q E_d(t)$ *has a non zero global section for* $1 \leq q \leq N\,,\ q+1 \leq \binom{n+t}{n}$ *and* $t \geq 1$.

Proof
The exact sequence $0 \longrightarrow \bigwedge^q E_d \longrightarrow \bigwedge^q S^dV \otimes \mathcal{O}_{\mathbb{P}^n} \longrightarrow \bigwedge^{q-1} E_d\,(d) \longrightarrow 0$ implies that

$$H^0\left(\bigwedge^q E_d(t)\right) \;=\; Ker\left(\bigwedge^q S^dV \otimes S^tV \xrightarrow{\alpha_t} \bigwedge^{q-1} S^dV \otimes S^{t+d}V\right).$$

Now, there is a Koszul complex

$$\longrightarrow \bigwedge^{q+1} S^dV \otimes \mathcal{O}(t-d) \longrightarrow \bigwedge^q S^dV \otimes \mathcal{O}(t) \xrightarrow{a_t} \bigwedge^{q-1} S^dV \otimes \mathcal{O}(t+d) \longrightarrow$$

with $\alpha_t = H^0(a_t)$. For $t \geq d$, global sections of $\bigwedge^{q+1} S^dV \otimes \mathcal{O}(t-d)$ will therefore give sections of $\bigwedge^q E_d(t)$. In particular, for $d = t$, we get that for each family $s_0, \cdots, s_q$ of degree d polynomials,

$$\sum_{i=0}^{q} (-1)^i \; s_0 \wedge \cdots \wedge \widehat{s_i} \wedge \cdots \wedge s_q \otimes s_i$$

is in the kernel of α_d. Let now be $1 \leq t < d$. If we can factor $s_i = u\, w_i$ with u of degree $d-t$, then

$$\sum_{i=0}^{q} (-1)^i \; s_0 \wedge \cdots \wedge \widehat{s_i} \wedge \cdots \wedge s_q \otimes w_i$$

must be in the kernel of α_t, and therefore defines a global section of $\bigwedge^q E_d(t)$. Thus to get a non-zero section of $\bigwedge^q E_d(t)$, it suffices to find $q+1$ linearly independent polynomials of degree t, which is possible as soon as $q+1 \leq \binom{n+t}{n}$. $\diamond$

Remark 2.3 The bundles $\bigwedge^q E_d$ are semistable (see [P], Proposition 5.6), so $H^0\left(\bigwedge^q E_d(t)\right) = 0$ if $\mu\left(\bigwedge^q E_d(t)\right) = t - \frac{qd}{N} < 0$. In particular,

$$H^0\left(\bigwedge^q E_d(t)\right) = 0 \quad \forall t \leq 0\,.$$

3 - Conclusions

In this section we will fit our results into the literature.
In particular, we will prove that:

Theorem 3.1 *Let d be an integer s.t. $d \geq 3$. Then the Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^2}(d)} : \mathbb{P}^2 \hookrightarrow \mathbb{P}^N$ satisfies property N_p if and only if $0 \leq p \leq 3d-3$. Moreover, if $d = 2$, the embedding $\varphi_{\mathcal{O}_{\mathbb{P}^2}(2)} : \mathbb{P}^2 \hookrightarrow \mathbb{P}^5$ satisfies N_p $\forall p$.*

We have the following:

Proposition 3.2 (M. Green, C. Birkenhake)
Let $d \geq 2$ and $p = \begin{cases} 3d-3 & \text{if } d \geq 3 \\ 2 & d = 2 \end{cases}$. Then the complete Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^2}(d)} : \mathbb{P}^2 \hookrightarrow \mathbb{P}^N$ satisfies property N_p.

Proof
See [B1], Corollary 3.2.
The result follows also applying Theorem 3.b.7 of [G1](which says that the minimal resolution of a Veronese variety restricts to the minimal resolution of its curve hyperplane section) and Theorem 4.a.1 of [G1] (which says that a line bundle of degree $2g+1+p$ on a curve of genus g satisfies N_p). $\diamond$

In the same way we get:

Lemma 3.3 *The Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^3}(3)} : \mathbb{P}^3 \hookrightarrow \mathbb{P}^{19}$ satisfies N_6.*

Proof
The curve hyperplane section of the image of the cubic Veronese embedding of $\mathbb{P}^3$ is the space curve complete intersection of two cubics embedded by $|\mathcal{O}_{\mathbb{P}^3}(3)|$ and it has genus 10. The result follows again applying Theorem 3.*b*.7 and Theorem 4.*a*.1 of [G1]. ◇

Lemma 3.4 *The ideal* $\mathcal{I}$ *of* $\varphi_{\mathcal{O}_{\mathbb{P}^2}(2)}(\mathbb{P}^2)$ *in* $\mathbb{P}^5$ *has the following resolution:*

$$O \longrightarrow \mathcal{O}_{\mathbb{P}^5}(-4)^{\oplus 3} \longrightarrow \mathcal{O}_{\mathbb{P}^5}(-3)^{\oplus 8} \longrightarrow \mathcal{O}_{\mathbb{P}^5}(-2)^{\oplus 6} \longrightarrow \mathcal{I} \longrightarrow 0\,.$$

In particular the Veronese embedding $\varphi_{\mathcal{O}_{\mathbb{P}^2}(2)}\colon \mathbb{P}^2 \hookrightarrow \mathbb{P}^5$ *satisfies* N_p $\forall p$.

Proof
Easy computation. ◇

Proof of Theorem 3.1
By Proposition 3.2 and Lemma 3.4 we just need to show that if $d \geq 3$ then property N_p does not hold for $p \geq 3d-2$. But this is exactly the bound coming from Theorem 2.1. ◇

When $d = 2$ the minimal free resolution of the quadratic Veronese variety is known from the work of Jozefiak, Pragacz and Weyman [JPW], in which they prove a conjecture made by Lascoux. As a corollary of the above paper we have the following result (which agrees with our Conjecture formulated in the Introduction):

Theorem 3.5 *The quadratic Veronese embedding* $\varphi_{\mathcal{O}_{\mathbb{P}^n}(2)} : \mathbb{P}^n \hookrightarrow \mathbb{P}^N$ *satisfies* N_p *if and only if* $p \leq 5$ *when* $n \geq 3$ *and* $\forall p$ *when* $n = 2$. ◇

The following nice characterization, probably well known, was found during discussions with E. Arrondo:

Theorem 3.6 *The only (smooth) varieties in* $\mathbb{P}^n$ *such that* N_p *holds for every* $p \geq 0$ *are the quadrics, the rational normal scrolls and the Veronese surface in* $\mathbb{P}^5$.

Proof
Suppose X is a variety satisfying N_p for every $p \geq 0$. Then $H^i(\mathcal{O}_X(t)) = 0$ for $t \geq 0$ and $1 \leq i \leq dim\, X - 1$. Hence from Theorem 3.*b*.7 in [G1] it follows that the minimal free resolution of X restricts to the minimal resolution of its generic curve section C. This implies that $H^1(\mathcal{O}_C) = 0$ and C is linearly normal, hence

C is a rational normal curve. In particular, X has minimal degree and we get the result. $\diamond$

We remark that the only Veronese varieties appearing in Theorem 3.6 are the rational normal curves and the Veronese surface in $\mathbb{P}^5$.

References

[**B1**] C. Birkenhake
Linear systems on projective spaces
Manuscripta Math. 88 (1995), 177-184

[**B2**] C. Birkenhake
Noncomplete linear systems on abelian varieties
Trans. A.M.S. vol 348, Number 5 (1996), 1885-1908

[**EL**] L. Ein, R. Lazarsfeld
Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension
Invent. Math. 111 (1993), 51-67

[**G1**] M. Green
Koszul cohomology and the geometry of projective varieties
J. Diff. Geometry 19 (1984), 125-171.

[**G2**] M. Green
Koszul cohomology and the geometry of projective varieties, II
J. Diff. Geometry 20 (1984), 279-289.

[**GL**] M.Green, R.Lazarsfeld
Some results on the syzygies of finite sets and algebraic curves
Compositio Math. 67 (1988), 301-314.

[**JPW**] T. Jozefiak, P. Pragacz, J. Weyman
Resolutions of determinantal varieties and tensor complexes associated with

symmetric and antisymmetric matrices
Asterisque 87-88 (1981), 109-189.

[**M**] L. Manivel
On the syzygies of flag manifolds
Proc. AMS 124 (1996), 2293-2299

[**P**] R. Paoletti
Stability of a class of homogeneous vector bundles on $\mathbb{P}^n$
Bollettino U.M.I. (7) 9-A (1995), 329-343

Authors' addresses

Giorgio Ottaviani
Dipartimento di Matematica
"U. Dini"
Università di Firenze
viale Morgagni 67/A
I 50134 FIRENZE
ottavian@math.unifi.it

Raffaella Paoletti
Dipartimento di Matematica ed
Appl. per l'Architettura
Università di Firenze
piazza Ghiberti
50122 Firenze
raffy@math.unifi.it